

---

**Please follow the black and white number and arrow signs for this tour.**

PLEASE TAKE A COPY OF THE GEOLOGIC TIME PERIODS SHEET WITH YOU. (Also in the plastic holder.)

To begin the tour, walk down the ramp, turn left and follow the path along the lake. You will see on both sides of the path several **(1) *Ginkgo biloba*** trees. *Ginkgo* is a “living fossil”, so-called because fossil records indicate the species has remained virtually unchanged for over 250 million years. (See the geologic time chart on the back.) The species name “*biloba*” refers to the double-lobed shape of their leaves. Ginkgos are dioecious, meaning they have separate male and female trees. Males produce pollen cones, while the females produce seed. The seeds have a rather unpleasant odour and thus female trees are not usually grown in public places. However we grow both male and female trees at VanDusen. As its long history attests, this is a very resistant tree. Disease, pollution, cold weather and even radioactivity do not kill it - a *Ginkgo biloba* growing about a kilometer from the epicentre, survived the atomic bombing of Hiroshima!

Further along the path is a trail off to the right (see arrow) into the Woodland Garden. Follow the path for a few feet and turn right at the next arrow onto the side path leading to a large **(2) coast redwood (*Sequoia sempervirens*)**. This species boasts the tallest tree in the world at 379 feet, found in Northern California in 2006. It was named ‘Hyperion’ by the team that discovered it. Coast redwoods can live for 2000 years, with trunks up to 21 feet in diameter. This one is just a baby! Its prehistoric ancestors thrived with the dinosaurs during the Jurassic Era, ~160 mya (million years ago).

Return to the path and turn right. You will come to a large **(3) southern magnolia (*Magnolia grandiflora*)**, native to southeastern USA, on your right. Magnolias exhibit what is called a disjunct distribution. They were once widespread throughout the world but, due to climate change, are now restricted to two widely separated regions - southeast Asia and southeastern North America and adjacent Central America. Magnolias evolved 60 million years ago, before bees evolved, and they are pollinated by beetles. This accounts for their very tough petals, resistant to crawling and chewing! This species produces large scented white flowers in spring. Beneath the magnolia, and in many other areas of the garden, are various kinds of ferns. This one is our native **(4) deer fern (*Blechnum spicant*)**. Ferns are amongst the oldest land plants, arising ~360 million years ago in the Carboniferous period. They are widespread and range in height from a few centimetres to several meters tall, the latter being tree ferns. Later in this tour you will see some tree ferns.

Continue along the woodland path towards Cypress Pond and turn left. On the other side of the pond you see a grove of deciduous **(5) bald cypress (*Taxodium distichum*)** trees, with feathery pale green foliage (which will not yet be visible in early April). These beautiful conifers, members of the pine family, were widespread on earth as far back as the Jurassic period ~150 mya. Continue along the main path and across the little stone bridge. Ahead on your right are **(6) the true cedars (*Cedrus* species)**, not to be confused with western redcedar, which isn’t a cedar at all but a species of *Thuja*. True cedars have needles in whorls of ~15 to 20. Three of the four species of *Cedrus* are represented at VanDusen. A useful way to distinguish them is to remember that *Cedrus atlantica* has ascending branches, cedar of Lebanon (*Cedrus libanii*) has level branches, and *Cedrus deodara* has descending branches. True cedars belong to the pine family and arose in the Tertiary period.

Turn left along the gravel path into the Southern Hemisphere Garden. On your right is a somewhat scraggly tree, **(7) Antarctic beech or ñire (*Nothofagus antarctica*)**. The genus *Nothofagus* has provided evidence to help prove the theory of continental drift. Today species of *Nothofagus* are found in South America, New Guinea, New Zealand and Australia. Pollen fossils have been found in Antarctica, indicating it must have existed on the super continent, Gondwana, between 180 and 510 mya. Continue along the path. Ahead of you are several **(8) Tasmanian tree ferns (*Dicksonia antarctica*)**. They are native to Australia and they can survive our winters if they are well wrapped to protect them from frost. Their ancestors date back to the Jurassic period. Turn left. Note on your left, just off the path, an enormous, beautiful, evergreen **coigüe (*Nothofagus dombeyi*)** from Chile and Argentina. This species often forms mixed forests with money puzzle trees. Its leafy evergreen boughs are used for thatching and its wood is decay resistant.

Ahead you will see several **(9) monkey puzzle trees, (*Araucaria araucana*)** also native to Chile and Argentina. The spiny leaves are extremely sharp! Ancestors of these trees lived 160 mya and, like ginkgos, the male and

female reproductive structures are on separate trees. At VanDusen we grow both sexes, distinguishable by the size and shape of their cones - the female cones are large and rounded while the male cones are smaller and more elongated.

Proceed down the path and across the little zig-zag bridge. In Livingstone Lake there are many water lilies. Most of the pond is covered with the yellow blossomed *Nuphar lutea*, found throughout the northern hemisphere. Their ancestors date back about 60 million years. Frogs and insects love them - humans not so much. Invasive, they clog the lake and must be removed regularly. On your right as you step off the bridge is a **(10) Japanese umbrella pine (*Sciadopitys verticillata*)**. It is not a pine at all and has no close relatives amongst present-day plants. It branched off from the other conifers ~250 mya, spreading throughout Eurasia, northern Europe and northern North America. Today it is restricted to a small area in Japan.

Proceed left along the path beside the lake. By the water's edge are several spectacular **(11) giant rhubarb plants (*Gunnera species*)**. These have one of the largest leaves of any plant in the world! Be careful if you touch them as the leaves are very rough. The flower stalks are enormous (you will find them underneath the leaves) but the tiny red flowers are inconspicuous. *Gunnera* species date back nearly 100 million years to the middle of the Cretaceous period. One of the fascinating things about these plants is that **cyanobacteria** live within their leaf cells. These single-celled organisms carry out photosynthesis and "fix" nitrogen, converting it from an inert gas to nitrate or ammonia, forms of nitrogen which plants can use. Because of this 'in house' nitrogen supply, *Gunnera* can grow in nitrogen-poor swampy areas. In turn, the cyanobacteria have a sheltered place to live. This is a nice example of symbiosis, similar to the way legumes obtain nitrogen from nitrogen-fixing bacteria living within their root nodules. An amazing thing about cyanobacteria is that their ancestors were amongst the first living things on Earth, dating back ~3.4 billion years! These ancestors colonized the leaf cells of early plants and evolved into chloroplasts, the organelles which carry out photosynthesis. Meanwhile, non-photosynthesizing bacteria colonized both plant and animal cells and evolved into mitochondria, the organelles which obtain energy from food. Without these, the advanced forms of life on Earth as we know them could not exist.

Proceed along the path beside the lake and watch for turtles sunning themselves on the rocks. At the end of the path turn right into the Phyllis Bentall Garden surrounding a rectangular pool. Across from the pool are pots containing various kinds of **(12) carnivorous plants**. These plants obtain nitrogen by trapping and digesting insects, and can therefore grow in nitrogen-poor soils. Also growing in these pots are miniature horsetail (*Equisetum scirpoides*), another living fossil. The genus *Equisetum* is even older than *Ginkgo*, dating back to the Devonian period ~400 mya. During the Carboniferous period some horsetail relatives were 65 feet tall, but they did not survive. Today horsetails are widely spread throughout the world, except for Australasia and Antarctica. Some are invasive and difficult to remove and gardeners consider them to be nuisance weeds, beautiful though they are.

This is the end of the tour. We hope you enjoyed your trip into the geologic past! To return to the garden entrance go back towards the path beside the lake and over the wooden bridge.

GEOLOGIC	MILLIONS OF	
TIME PERIODS	YEARS AGO	
- not to scale -	(mya)	ORGANISMS:

---

Quaternary		
-----1.6		Humans (~4 mya)
Tertiary		<b>Magnolia, Cedrus, Nuphar, Veronica, Gunnera</b>
-----66		<b>Equisetum</b> (~65mya)
Cretaceous		<b>Nothofagus</b> (~80 mya),
		<b>Gunnera</b> (~95 mya)
-----138		First flowering plants
Jurassic		<b>Taxodium</b> (Bald Cypress), <b>Dicksonia</b> (tree ferns), Wollemi
		pine
		Monkey puzzle tree
-----205		<b>Ginkgo biloba</b> (~200 mya, living fossil)
Triassic		First dinosaurs (~230 mya)
-----240		
Permian		<b>Sciadopitys</b> (~250 mya)
-----290		First conifers
Carboniferous		
-----360		First ferns
Devonian		
-----410		Early horsetails (~400 mya) (Coelacanth fishes ~390 mya -
		living fossils)
Silurian		
-----435		
Ordovician		First land plants (~470 mya) - mosses, liverworts
-----500		
Cambrian		First marine animals - e.g. sponges, earliest fishes
-----545		
Pre-Cambrian		
		First fungi (~1300 mya)
		<b>Cyanobacteria</b> - among first living things (~3400 mya) -
		photosynthesizers
-----3800		
Hadean		
Formation of earth -----4600		